

HIV Medication Chart

Nucleoside/Nucleotide Analogue Reverse Transcriptase Inhibitors (NRTI)

Emtriva*
(emtricitabine, FTC)



Epivir*†
(lamivudine, 3TC)



Retrovir*†
(zidovudine, AZT, ZDV)



Viread*
(tenofovir, TDF)



Ziagen*†
(abacavir, ABC)



Combination Antiretrovirals

Atripla
(EFV+FTC+TDF)



Combivir †
(3TC+AZT)




Complera
(FTC+RPV+TDF)



FDA Pregnancy Category D

Descovy
(FTC+TAF)



Epzicom ▲
(ABC+3TC)




Evotaz
(ATV+COBI)



Genvoya

(EVG+COBI+FTC+TAF)



Odefsey

(FTC+RPV+TAF)



Prezcobix


(DRV+COBI)



Stribild
(EVG+COBI+FTC+TDF)




Triumeq
(ABC+DTG+3TC)



Trizivir †
(ABC+3TC+AZT)



Truvada
(FTC+TDF)



Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTI)

Edurant
(rilpivirine, RPV)



Intellece
(etravirine, ETV)



Sustiva (efavirenz, EFV)



FDA Pregnancy Category D

Viramune*†
(nevirapine, NVP)



Protease Inhibitors (PI)

Kaletra*
(lopinavir/ritonavir, LPV/r)



Lexiva*
(fosamprenavir, FPV)



Prezista*
(darunavir, DRV)



Reyataz*
(atazanavir, ATV)



Viracept*
(nelfinavir, NFV)



Entry Inhibitors

Fuzeon
(enfuvirtide, T-20)
Fusion Inhibitor



Selzentry
(maraviroc, MVC)
CCR5 Antagonist



Boosting Agents

Norvir*
(ritonavir, RTV)



Tyboost
(cobicistat, COBI)



Integrase Inhibitors

Isentress▲*
(raltegravir, RAL)









Tivicay
(dolutegravir, DTG)



All pills shown in actual size. Medication brand names appear in bold. Generic names and commonly used abbreviations appear in parentheses. Bioavailability and indications between formulations may be different. * Also available in liquid or powder form. † Generic formulation available. ▲ Chewable form available.

Medication Schedule

Name _____ Date _____

Name of Medication	Number of pills to take each time	Time of day you are to take this medicine	Food Interactions	Side Effects	TOTAL number of pills each day
					
					
					
					
					
					

Helpful Hints:

- Refill prescriptions before you run out. Call for refills when you have at least 3 or 4 days left.
- Use cues as a reminder to take your pills (after a meal or favorite TV show, or before bedtime).
- Use reminder aids such as beepers, alarm clocks, and pillboxes. Ask your pharmacist about these.
- Plan ahead (vacations, travel, count out weekly doses).
- Do not stop taking your medications until you have spoken with your health care provider or pharmacist.
- If you have a severe reaction or in case of emergency, contact your health care provider IMMEDIATELY.

Contact Information: Provider _____

Clinic Phone _____ Pharmacy _____ Phone _____



Colorado AIDS Education & Training Center
University of Colorado, Anschutz Medical Campus
12631 E. 17th Ave., MS A089, Aurora, CO 80045

Tel: 303-724-0646 Fax 303-724-7297 www.caetc.org

Developed by Lisa Lawrence, MSW; and Steven Johnson, MD

Reviewers: Jasjit Gill, PharmD, University of Colorado Hospital ID Pharmacy; Jacob Langness, PharmD, University of Colorado Hospital ID Pharmacy; Judith Bagley, RPh, Denver Health Medical Center ID Pharmacy

Infrequently Used

Protease Inhibitors (PI)

Aptivus*
(tipranavir, TPV)



Crixivan
(indinavir, IDV)




Invirase
(saquinavir hard gel capsules, SQV)



Nucleoside/Nucleotide Analogue Reverse Transcriptase Inhibitors (NRTI)

Videx EC*†
(didanosine, ddl)



Zerit*†
(stavudine, d4T)



Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTI)

Rescriptor
(delavirdine, DLV)



Integrase Inhibitors

Vitekta*
(elvitegravir, EVG)



Discontinued Medications or Formulations

Agenerase
(amprenavir, APV)



Fortovase
(saquinavir soft gel capsule, SQV)



Hivid
(zalcitabine, ddC)



Kaletra
(lopinavir/ritonavir, LPV/r)



Videx
(didanosine, ddl)

