


Sofosbuvir/Velpatasvir (Epclusa®) for Genotype 3 No Cirrhosis*

Preliminary data suggest that NS5A Genotype 3 RAV testing is recommended for all genotype 3 patients prior to treatment. Consult Specialist first**.



*See next page for Epclusa decision tree in persons with cirrhosis.

**Consult Liver Disease Specialist before prescribing treatment for persons with genotype 3, past treatment failure, cirrhosis, hepatocellular carcinoma (HCC), renal disease, pre- and post-liver transplant, or HIV or hepatitis B co-infection.


Link to AASLD/IDSA Guidelines: <http://hcvguidelines.org/full-report-view>

¹Peg/RBV = Pegylated interferon/
Ribavirin

²Sof/RBV = Sofosbuvir/Ribavirin

Sofosbuvir/Velpatasvir (Epclusa®) for Genotype 3 Cirrhosis*

Preliminary data suggest that NS5A Genotype 3 RAV testing is recommended for all genotype 3 patients prior to treatment. Consult Liver Disease Specialist first*.



*Consult Liver Disease Specialist before prescribing treatment for persons with genotype 3, past treatment failure, cirrhosis, hepatocellular carcinoma (HCC), renal disease, pre- and post-liver transplant, or HIV or hepatitis B co-infection.

**Epclusa 24 weeks- Consider for persons with decompensated cirrhosis who cannot take ribavirin.

Link to AASLD/IDSA Guidelines: <http://hcvguidelines.org/full-report-view>

¹Peg/RBV = Pegylated interferon/
Ribavirin

²Sof/RBV = Sofosbuvir/Ribavirin