

Viekira Pak™ (Ombitasvir/Paritaprevir/Ritonavir; Dasabuvir) or Viekira XR™ (Dasabuvir/Ombitasvir/Paritaprevir/Ritonavir) & Ribavirin Treatment Agreement

Liver Disease & Hepatitis Program Providers: Brian McMahon, MD; Youssef Barbour, MD; Lisa Townshend-Bulson, FNP-C; Annette Hewitt, FNP-C; Prabhu Gounder, MD; Ellen Provost, DO; Timothy Thomas, MD; Stephen Livingston, MD

Family Medicine Provider: _____

If you are considering hepatitis C treatment, please read this treatment agreement carefully and be sure to ask any questions you may have before you sign the form.

The FDA approved ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets co-packaged (Viekira Pak™) and dasabuvir, ombitasvir, paritaprevir, and ritonavir extended-release tablets (Viekira XR™) to be given with ribavirin for the treatment of hepatitis C genotype 1a without cirrhosis or with compensated cirrhosis (Child-Pugh Class A), including those with HCV/HIV co-infection, as well as genotype 1 liver transplant recipients with normal hepatic function and mild fibrosis.

Treatment with Viekira Pak™ or Viekira XR™ requires 6 scheduled visits over 6 months if your treatment course is 12 weeks and 9 scheduled visits over 9 months if your treatment course is 24 weeks.

PREGNANCY & BREASTFEEDING WARNING

It is not known if Viekira Pak™ or Viekira XR™ will harm an unborn or breastfeeding baby. However, ribavirin can harm an unborn child or breastfeeding infant. A woman must not get pregnant and a man must not father a child while taking ribavirin or for 6 months after treatment. You must **use 2 forms of birth control** when you are taking ribavirin and for 6 months after your last dose.

You must stop using ethinyl estradiol-containing medicines before you start treatment with Viekira Pak™ or Viekira XR™. If you use these medicines as a method of birth control you must use another method of birth control during treatment with Viekira Pak™ or Viekira XR™, and for about 2 weeks after finishing treatment with Viekira Pak™ or Viekira XR™.

Acceptable Birth Control Methods (must use 2):

Progestin only contraceptives (e.g. mini pill, Depo shot, Nexplanon™)

Male or female condom

Spermicides (creams, films, foams, gels, and/or suppositories)

Diaphragm or cervical cap

Intrauterine device (IUD), Today™ vaginal sponge

Unacceptable Birth Control Methods:

Rhythm method or withdrawal

HOW THE TREATMENT PROCESS WORKS

You will have blood and urine tests.

- These tests will include a pregnancy test for female patients of childbearing age. Urine pregnancy tests will be done monthly during clinic visits. If you are a woman and your treatment includes ribavirin it is recommended that you continue monthly home pregnancy testing for 6 months after treatment and notify your healthcare provider if you become pregnant. Female partners of males whose treatment includes ribavirin should do a monthly home pregnancy test during treatment and for 6 months after treatment completion and notify their health care provider if they become pregnant.
- Random drug and alcohol tests may be requested.
- At each visit, about 2-3 tubes of blood will be collected. Other examinations and tests may be done during the treatment if your provider feels there is a need.

Liver Clinic Provider, select the appropriate treatment regimen:

___ Viekira Pak™ or Viekira XR™ & ribavirin for 12 weeks for genotype 1a without cirrhosis

___ Viekira Pak™ or Viekira XR™ & ribavirin for 24 weeks for genotype 1a with compensated cirrhosis

Ribavirin dose is based on weight (1000 mg for persons less than or equal to 75 kg (165 lb.) and 1200 mg/day for those greater than 75 kg, divided and taken twice daily with food) except for those post liver transplant, see below.

Note: Viekira Pak™ and Viekira XR™ are contraindicated in persons with moderate to severe cirrhosis (Child-Pugh Class B or C).

Your first three visits will be at the start of treatment (week 0) and weeks 2 and 4 after you begin taking the medications. After that, the visits will be once each month until you stop taking the medications.

You may need to come to the clinic or see your primary care provider more frequently if you are having side effects or problems related to the treatment.

You will have a liver clinic visit 3 months after treatment completion and then yearly (corresponding to your end of treatment date) for 5 years. If you have cirrhosis you should continue to have a liver ultrasound and alpha fetoprotein (AFP) cancer screening blood test every six months and regular clinic visits.

TREATMENT MEDICATIONS AND SIDE EFFECTS

Viekira Pak™ contains ombitasvir, paritaprevir, and ritonavir tablets and dasabuvir tablets co-packaged for oral use. Two pink tablets contain ombitasvir 12.5mg, paritaprevir 75mg and ritonavir 50mg and are taken at the same time daily (in the morning) with a meal. Two beige tablets contain dasabuvir 250 mg and one of them is taken twice daily (in the morning and evening) with a meal. Store the medication at room temperature.

If you miss a dose of the pink tablets, and it is less than 12 hours from the time you usually take your dose, take the missed dose with a meal as soon as possible. Take your next dose at your regular time the next day with a meal. If you miss a dose of the pink tablets more than 12 hours from the time you usually take your dose, do not take the missed dose. Take your next dose at your regular time the next day with a meal.

If you miss a dose of the beige tablet and it is less than 6 hours from the time you usually take your dose, take the dose with a meal as soon as possible. Then take your next dose at your regular time with a meal. If it is more than 6 hours since you missed your dose, do not take the missed dose. Instead, take your next dose at your regular time with a meal.

Viekira XR™ contains dasabuvir, ombitasvir, paritaprevir, and ritonavir extended-release tablets, for oral use. Three pale yellow-colored tablets contain dasabuvir 200mg, ombitasvir 8.33mg, paritaprevir 50mg, and ritonavir 33.33mg and are taken at the same time daily with a

meal. Taking the tablets in a fasting state could result in reduced cure and the development of resistance. Swallow the tablets whole and do not consume alcohol within 4 hours of taking Viekira XR. Do not skip or miss doses. Store the medication at or below 86°F.

Do not take more than the prescribed dose of Viekira Pak™ or Viekira XR™ (no doubling up).

Do not take Viekira Pak™ or Viekira XR™ if you have had a severe skin rash after taking ritonavir (Norvir®).

The most common side effects of Viekira Pak™ or Viekira XR™ are nausea, itching and sleep problems.

Tell your healthcare provider if you are taking any of the following medicines; as they are contraindicated with Viekira Pak™ or Viekira XR™:

- Alfuzosin hydrochloride (Uroxatral™)
- Colchicine (COLCRYS, Mitigare®)
- Carbamazepine (Carbatrol™, Epitol™, Equetro™, Tegretol™); Phenytoin (Dilantin™, Phenytek™); Phenobarbital (Luminal™); Primidone (Mysoline®)
- Dronedarone (Multaq®); Lurasidone (Latuda®)
- Efavirenz (Atripla™, Sustiva™)
- Ergot containing medicines including: ergotamine tartrate (Cafergot™, Ergomar™, Ergostat™, Medihaler™); dihydroergotamine mesylate (D.H.E. 45™, Migranal™); methylergonovine (Methergine®); ergonovine (Ergotrate®)
- Ethinyl estradiol-containing medications; combination birth control pills or patches, such as Lo Loestrin™ FE, Norinyl™, Ortho Tri-Cyclen Lo™, Ortho Evra™; hormonal vaginal rings such as NuvaRing®; hormonal replacement therapy medicine Fem HRT™.
- Gemfibrozil (Lopid™); Lovastatin (Advicor™, Altoprev™, Mevacor™); Simvastatin (Simcor™, Vytorin™, Zocor™)
- Midazolam, when taken by mouth; Triazolam (Halcion™)
- Pimozide (Orap™)
- Ranolazine (Ranexa®)
- Rifabutin (Mycobutin®); Rifampin (Rifadin™, Rifamate™, Rifater™, Rimactane™)

- Sildenafil citrate (Revatio™) when taken for pulmonary artery hypertension
- St. John's wort (Hypericum perforatum) or a product that contains St. John's wort

Tell your healthcare provider if you are taking any of the following medicines; as they are not recommended with Viekira Pak™ Viekira XR™:

- Darunavir (Prezista®) / ritonavir; lopinavir/ ritonavir (Kaletra®); rilpivirine (Edurant®, Complera®, Odefsey®)
- Salmeterol (Serevent, Advair®)

Tell your healthcare provider if you are taking any of the following medicines; as dosage adjustments or monitoring may be recommended:

- Amiodarone (Cordarone®, Nexterone®, Pacerone®); bepridil; disopyramide (Norpace®, Norpace CR®); flecainide (Tambocor™); systemic lidocaine (Xylocaine®); mexiletine; propafenone (Rythmol, Rythmol SR); quinidine (Nuedexta®)
- Valsartan (Diovan®, Exforge®, Entresto™); losartan (Cozaar®, Hyzaar®); candesartan (Atacand®)
- Amlodipine (Norvasc®); Nifedipine (Procardia®, Adalat®); Diltiazem (Cardizem®, Tiazac®); Verapamil (Covera-HS®, Calan®, Verelan®)
- Furosemide (Lasix®)
- Ketoconazole; Voriconazole (Vfend®)
- Fluticasone (Inhaled- Arnuity™ Ellipta®, Breo Ellipta®, Flovent®, Advair®; Nasal – Flonase®, Veramyst, Dymista®)
- Atazanavir (Reyataz®)/ ritonavir (when co-administered with Viekira Pak, atazanavir 300mg without ritonavir should only be given in the morning)
- Cyclosporine (Gengraf®, Neoral®, Sandimmune®); tacrolimus (Astagraf XL®, Envarsus XR™, FK506 (common name), Hecoria™, Prograf®)
- Buprenorphine/naloxone (Suboxone®)
- Omeprazole (Prilosec®, Prilosec OTC®) (avoid use of > 40mg daily)
- Alprazolam (Xanax®)
- Quetiapine (Seroquel®)

- Rosuvastatin (Crestor®) dose not to exceed 10mg daily; pravastatin (Pravachol®) dose not to exceed 40mg daily

Ribavirin is a 200mg capsule or tablet. You will take ribavirin pills twice daily by mouth with food (dose is based on your weight, except for those who have had a liver transplant). Ribavirin dose may be adjusted based on your tolerance of this medication. You should not miss/skip taking any pills. A common side effect is anemia. Anemia is a condition where the blood has a decreased number of red blood cells. This occurs more often in older persons taking ribavirin. Anemia can be serious in patients who have kidney problems. In patients who have coronary artery disease (narrowing of the blood vessels in the heart), anemia may make the problem worse, leading to chest pain or heart attack. If your provider believes you may have coronary artery disease, you will be tested for this and excluded from treatment if it is serious.

- Other common side effects include: headache, trouble sleeping, nausea, vomiting, weakness or lack of energy, shortness of breath, loss of appetite, itching, cough, muscle pain, swelling and pain in your joints (gout), depression, nervousness, and dizziness.
- Studies in animals have shown when ribavirin is given to pregnant females, death of the developing embryo or birth of deformed baby animals may result. It is expected that similar results as seen in the animal studies could occur in humans.

The most common side effects of Viekira Pak™ or Viekira XR™ when given with ribavirin are tiredness, nausea, itching, skin reactions such as redness or rash, sleep problems, and feeling weak.

PLEASE NOTE:

You must let your medical, mental health, dental providers, and pharmacist(s) know that you are taking Viekira Pak™ or Viekira XR™ & ribavirin prior to starting any new medications. You must let Liver Clinic providers know about any new medications you are prescribed before starting them. This includes vitamins and other supplements.

*** Hepatitis C treatment should not cause pain that requires narcotic pain medication.

BENEFITS OF TREATMENT

In most cases, hepatitis C will respond to treatment as determined by a blood test that measures the presence and amount of hepatitis C in the blood. If you have no hepatitis C in your blood 12 weeks **after** the end of treatment, this is called a “sustained virologic response” and means you no longer have hepatitis C. Your chance of achieving a sustained virologic response depends on the hepatitis C genotype, how much hepatitis C virus you have in your blood at the beginning of treatment, any past treatment response, and how much liver damage you have had prior to treatment.

It is possible that you may develop some serious side effects, which will require you to stop the treatment. You may still benefit from treatment even if it does not get rid of your hepatitis C, as it may slow down the disease. You may choose to stop treatment at any time.

In Clinical Trials:

Persons with genotype 1a who did not have cirrhosis were treated with Viekira Pak™ (the components of Viekira XR™) and ribavirin for 12 weeks and had a 94% or better response (cure) rate.

Those with genotype 1a with compensated cirrhosis treated with Viekira Pak™ and ribavirin for 24 weeks had a 95% response rate.

Liver transplant recipients who were 12 months or longer after transplant and had normal liver function and mild fibrosis were given Viekira Pak™ (the components of Viekira XR™) and ribavirin for 24 weeks. Of 34 subjects, 97% (28/29) of those who were genotype 1a and 100% (5/5) of those who were genotype 1b responded.

WHOM TO CALL

If you have any questions about your treatment, contact the Liver Disease & Hepatitis Program @ 907-729-1560 or your primary care provider.

TREATMENT AGREEMENT

To receive treatment, please review the following statements and initial beside the responses:

_____ I agree not to drink alcohol or use recreational drugs during the treatment.

_____ I will tell my provider if I have any serious medical conditions (such as heart disease, high blood pressure, diabetes, high cholesterol, rheumatoid arthritis, or drug addiction), or psychiatric conditions (depression, history of suicide attempts, bipolar disorder, or psychosis).

_____ I am willing to visit the clinic and see a provider on a regular schedule for the entire length of the treatment. If I am unable to attend an appointment, I will let my provider know this ahead of time and I will reschedule my appointment.

_____ I understand that my treatment will be stopped if I cannot attend appointments as required to evaluate my health and well-being during treatment and the effectiveness of treatment.

_____ I will use 2 acceptable methods of birth control during treatment and for 6 months after I stop treatment (see lists, page 2).

_____ As a female, I understand that I cannot be pregnant or breastfeeding during the treatment and for 6 months after treatment. I understand that my treatment will be stopped if I become pregnant. _____ Not applicable, I am surgically sterile or post-menopausal.

_____ As a male taking ribavirin I understand that I should not father a child during treatment and for 6 months after treatment.

_____ If I have any problems with the medications or side effects that bother me, I will let my provider or nurse know right away.

_____ I understand that my hepatitis C may not respond to treatment.

_____ I understand that my provider can stop my treatment if the provider feels that stopping it is in the best interest of my health and welfare.

_____ I will do my best to take my medications as prescribed by my provider. If I am unable to do so, I will contact my provider.

_____ I will protect myself and others from hepatitis C by not sharing needles, toothbrushes, razors or nail clippers and covering cuts to prevent blood exposure.

My signature below means that I have read this treatment agreement and/or the meaning of the information has been explained to me. I agree to treatment.

Patient's Name (PLEASE PRINT)

Patient's Signature

Date

Provider's Name (PLEASE PRINT)

Provider's Signature

Date