

Hepatitis C Health Summary

Name: _____

DOB: _____

Phone #: _____

Alternate Contact: _____

Medications²:

Allergies:

Labs Prior to Treatment:

- Immediately prior: Pregnancy test
 - Uric Acid (ribavirin only)
- Within 1 month: CBC with differential
 - CMP (If GFR <30, do not start tx ¹)
 - PT/INR
 - HCV RNA
- Within 3 months: Genotype confirmation
 - HBV DNA (if HBV cAb or sAg +)
- Within 6 months: AFP
 - TSH
 - A1C or Fasting Glucose
 - Vitamin D 25OH
- Within 1 year: HIV screening
 - NS5A RAV (genotype 3 only)
- Once: IL-28b (if considering 8 weeks)

Pertinent Medical History:

- Previous hepatitis C treatment¹ Yes No
Specify: _____
- Cirrhosis¹ Yes No
Child-Pugh Score: _____
- Other Liver Disease¹ Yes No
Specify: _____
- Pulmonary Disorders¹ Yes No
Specify: _____
- Cardiac Disease² Yes No
Specify: _____
- DVT or PE¹ Yes No
Specify: _____
- PPI/H2 blocker/Antacid use² Yes No
Specify: _____
- Autoimmune Disorders² Yes No
Specify: _____
- Cancer Yes No
Specify: _____
- Current infection¹ Yes No
Specify: _____
- High Blood Pressure Yes No
- High Cholesterol Yes No
- Kidney Disease² Yes No
- Anemia^{1,2} Yes No
- Current TB Treatment² Yes No
- Diabetes Specify Type 1 or 2 Yes No
- HIV or AIDS¹ Yes No
- Seizure Disorder² Yes No
- Depression/Anxiety Yes No
- Other Psychiatric Conditions Yes No
Specify: _____

Screen & Review: AUDIT-C ___ PHQ-9 ___
 Vaccine Status (give if needed):
 Hepatitis A ___ (If unknown, check hep A total IgG)
 Hepatitis B ___ (If unknown, check HBsAg & HBsAb)

- Other vaccines as appropriate:
- Flu (annually)
 - PCV-13 (≥ age 65 or immunosuppressed)
 - PPSV-23 (≥ age 50 AN/AI in AK or high risk)
 - Td (once every 10 years) **OR** Tdap (once)
 - Zoster (≥ age 60)
 - ECG (over age 65 or h/o cardiac disease)

Birth Control: Birth Control Methods: _____
 Females: LMP: _____ Pregnant Yes No
 Males: Is your partner pregnant? Yes No
 Counsel about pregnancy prevention (see Treatment Agreement)
 Hepatitis C Treatment Agreement reviewed and signed

1- Further evaluation as indicated; consult Liver Disease Specialist prior to treatment.
 2- Check drug interactions to treatment drugs. Further evaluation as indicated.



ALASKA NATIVE TRIBAL HEALTH CONSORTIUM

Liver Disease & Hepatitis Program
4315 Diplomacy Drive, Anchorage, AK 99508
Phone: 907-729-1560 Fax: 907-729-1570

<http://www.anthc.org/hep>

Follow us on Twitter:

Liver Program @ANTHCLiver

We are glad to hear you are interested in treatment for hepatitis C!

Here are some things to think about (and do) before you make your final decision about treatment:

Why do treatment now? New medicines have increased the chance of cure and have fewer side effects.

Some people have worse liver disease than others. If you have more severe liver disease (a lot of scarring in the liver or cirrhosis) you should consider getting treatment sooner.

What will happen during treatment?

There are 6 FDA approved treatment options for **genotype 1**:

- Option 1 is Harvoni[®] (ledipasvir/sofosbuvir), 1 tablet taken once a day for 8-24 weeks. The most common side effects are feeling tired and headache. In clinical studies, treatment response rates to Harvoni[®] were 94-100%.
- Option 2 is Epclusa[®] (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The most common side effects are headache and feeling tired. In clinical studies, treatment response rates to Epclusa[®] were 94-98% for genotype 1.
- Option 3 is Viekira Pak[™] (ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets) co-packaged as 3 tablets in the morning and 1 tablet in the evening with food for 12-24 weeks. The major side effects are nausea, itching, and insomnia. In clinical studies, response rates to Viekira Pak[™] treatments were 86-100%.
- Option 4 is Zepatier[™] (elbasvir/grazoprevir), 1 tablet taken once a day for 12-16 weeks. The most common side effects are feeling tired, nausea, and headache. In clinical studies, treatment response rates to Zepatier[™] were 95-100%.
- Option 5 is Olysio[®] (simeprevir) plus Sovaldi[®] (sofosbuvir) 1 tablet of each taken once daily for 12 weeks. The most common side effects are feeling tired, headache, and nausea. In clinical studies, treatment response rates to Olysio[®] and Sovaldi[®] were 86-100%.
- Option 6 is Daklinza[™] (daclatasvir) plus Sovaldi[®] (sofosbuvir) 1 tablet of each taken once daily for 12 weeks. The most common side effects are headache and feeling tired. In clinical studies, treatment response rates to Daklinza[™] and Sovaldi[®] were 50-100%.

The FDA-approved **Genotype 2** treatment is Epclusa[®] (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The major side effects are headache and feeling tired. In clinical studies, the treatment response rate to Epclusa[®] was 99% for genotype 2.

There are 2 FDA-approved treatment options for **genotype 3**:

- Option 1 is Eplusa® (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The most common side effects are headache and feeling tired. In clinical studies, treatment response rates to Eplusa® were 85-98% for genotype 3.
- Option 2 is Daklinza™ (daclatasvir) and Sovaldi® (sofosbuvir) 1 tablet of each taken once daily for 12 weeks. The most common side effects are headache and feeling tired. In clinical studies, treatment response rates for Daklinza™ and Sovaldi® were 58-98%.

Some treatments will require ribavirin which is 5-6 additional tablets divided between morning and evening with food. The major side effects are feeling tired, nausea, itching and skin rash, trouble sleeping, irritability and weakness. A common side effect of ribavirin is anemia.

PLEASE NOTE: Ribavirin cannot be given to a pregnant or breastfeeding female or to a female who plans to become pregnant or a male who plans to father a child during or for 6 months after treatment because it can cause birth defects. There are no studies on Harvoni®, Eplusa®, Sovaldi®, Viekira Pak™, Zepatier™, or Daklinza™ in pregnant women or nursing mothers. Safety/risk during pregnancy or breastfeeding has not been established.

Are you ready for treatment?

To ensure that you will be successful in completing hepatitis C treatment we ask that the following items be done before starting treatment. We will review them together.

- You must be alcohol and drug-free. If you have recent drug/alcohol abuse, you need to be in an approved drug treatment program.
- You need to discuss hepatitis C treatment with your primary care provider and get his or her "OK" to start treatment.
- You should have a relative/close friend who is willing to help support you during treatment.
- You need to be committed to making every treatment appointment and getting **FREQUENT** blood draws (every 1-4 weeks). We will want to follow you very closely during treatment.

Additional Requirements If Checked:

_____ If you have cirrhosis, you may need an EGD (when a doctor looks into your esophagus and stomach for swollen veins that can bleed).

_____ If you have cirrhosis, you need to have an ultrasound of the liver (done in the past 6 months). This ultrasound checks your liver for cancer.

Once everything you need to do on the list has been done, call your primary care provider to make an appointment to plan for hepatitis C treatment. At this appointment, treatment and side effects will be discussed in detail.

If you are coming to Anchorage and want a Fibroscan, call the Liver Clinic ahead of your visit to schedule. Fibroscan is a test using ultrasound waves to check liver stiffness or scarring/fibrosis in your liver. Fibroscan testing is done in the Internal Medicine Clinic. Do not eat or drink for 3 hours before the test.

Congratulations on completing all the pre-treatment requirements!

Hepatitis C Treatment Checklists

Prior to Treatment

Labs

- | | |
|--|--|
| Immediately prior: ___ Pregnancy test (if applicable)
___ Uric Acid (with ribavirin)
Within 1 month: ___ CBC with differential
___ CMP ¹
___ PT/INR
___ HCV RNA
Within 3 months: ___ Genotype confirmation
___ HBV DNA (if HBV cAb or sAg +)
Within 6 months: ___ AFP
___ TSH
___ A1C or Fasting Glucose
___ Vitamin D 25OH (treat if deficient)
Within 1 year: ___ HIV screening
___ NS5A RAV (genotype 3 only)
Once: ___ IL-28b (if considering 8 weeks) | Miscellaneous:
___ Hepatitis A (If vaccine status is unknown, draw HAV total)
___ Hepatitis B (If vaccine status is unknown, draw HBsAg & HBsAb)
___ PHQ-9 baseline
___ AUDIT-C
___ Counsel about pregnancy prevention
___ Review & sign Treatment Agreement |
|--|--|

8 week

Week 4

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Week 8

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

12 week

Week 2 (with ribavirin)

- ___ CBC
- ___ CMP¹

Week 4

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Week 8

- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Week 12

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

16 week

Week 2 (with ribavirin)

- ___ CBC
- ___ CMP¹

Week 4

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Weeks 8 & 12

- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Week 16

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

24 week

Week 2 (with ribavirin)

- ___ CBC
- ___ CMP¹

Week 4

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Weeks 8, 12, 16, & 20

- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Week 24

- ___ HCV RNA
- ___ CBC
- ___ CMP¹
- ___ Pregnancy test

Nurse follow-up in clinic or by phone:

- ___ Managing side effects
- ___ Medication adherence discussion
- ___ Alcohol intake
- ___ Birth control reminder
- ___ Refill reminder

3 months post treatment

- ___ CBC
- ___ Liver Function Tests
- ___ HCV RNA
- ___ AUDIT-C

6 months post treatment

- ___ HCV RNA
- ___ AFP
- ___ RUQ US (if advanced fibrosis)
- ___ AUDIT-C

1- Sofosbuvir- or daclatasvir-based regimen - If GFR <30, no safe recommendation.
With ribavirin - If GFR <50, decrease dose (refer to package insert).

Viekira Pak™ (Ombitasvir/Paritaprevir/Ritonavir; Dasabuvir) or Viekira XR™ (Dasabuvir/Ombitasvir/Paritaprevir/Ritonavir) Treatment Agreement

Family Medicine Provider: _____

If you are considering hepatitis C treatment, please read this treatment agreement carefully and be sure to ask any questions you may have before you sign the form.

The FDA approved ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets co-packaged (Viekira Pak™) and dasabuvir, ombitasvir, paritaprevir, and ritonavir extended-release tablets (Viekira XR™) for the treatment of hepatitis C genotype 1b, including those with HCV/HIV co-infection.

Treatment with Viekira Pak™ or Viekira XR™ requires 6 scheduled visits over 6 months for a treatment course of 12 weeks.

PREGNANCY & BREASTFEEDING WARNING

It is not known if Viekira Pak™ or Viekira XR™ will harm an unborn or breastfeeding baby, so it is recommended that women do not get pregnant or breastfeed while taking this medicine.

You must stop using ethinyl estradiol-containing medicines before you start treatment with Viekira Pak™ or Viekira XR™. If you use these medicines as a method of birth control you must use another method of birth control during treatment with Viekira Pak™ or Viekira XR™, and for about 2 weeks after finishing treatment with Viekira Pak™ or Viekira XR™. Progestin only contraceptives (e.g. mini pill, Depo shot, Nexplanon™) are safe to use during treatment with Viekira Pak™ or Viekira XR™.

HOW THE TREATMENT PROCESS WORKS

You will have blood and urine tests.

- These tests will include a pregnancy test for female patients. A urine pregnancy test will be done monthly during a clinic visit.
- Random drug and alcohol tests may be requested.
- At each visit, about 2-3 tubes of blood will be collected. Other examinations and tests may be done during the treatment if your provider feels there is a need.

Provider, choose appropriate treatment regimen:

Viekira Pak™ or Viekira XR™ will be given for 12 weeks if:

____ You have genotype 1b and do not have cirrhosis.

____ You have genotype 1b and have mild cirrhosis. (Contraindicated in persons with moderate to severe cirrhosis)

Your first visit will be at the start of treatment (week 0) and then visits will be once each month until you stop taking the medications.

You may need to see your primary care provider more frequently if you are having side effects or problems related to the treatment.

You will have a clinic visit 3 months after treatment completion and then yearly (corresponding to your end of treatment date) for 5 years. If you have cirrhosis you should continue to have a liver ultrasound and alpha fetoprotein (AFP) cancer screening blood test every six months and regular clinic visits.

TREATMENT MEDICATIONS AND SIDE EFFECTS

Viekira Pak™ contains ombitasvir, paritaprevir, and ritonavir tablets and dasabuvir tablets co-packaged for oral use. Two pink tablets contain ombitasvir 12.5mg, paritaprevir 75mg, and ritonavir 50mg and are taken at the same time daily (in the morning) with a meal. Two beige tablets contain dasabuvir 250mg and one of them is taken twice daily (in the morning and evening) with a meal. Store the medication at room temperature.

If you miss a dose of the pink tablets, and it is less than 12 hours from the time you usually take your dose, take the missed dose with a meal as soon as possible. Take your next dose at your regular time the next day with a meal. If you miss a dose of the pink tablets more than 12 hours from the time you usually take your dose, do not take the missed dose. Take your next dose at your regular time the next day with a meal.

If you miss a dose of the beige tablet and it is less than 6 hours from the time you usually take your dose, take the dose with a meal as soon as possible. Then take your next dose at your

regular time with a meal. If it is more than 6 hours since you missed your dose, do not take the missed dose. Instead, take your next dose at your regular time with a meal.

Viekira XR™ contains dasabuvir, ombitasvir, paritaprevir, and ritonavir extended-release tablets, for oral use. Three pale yellow-colored tablets contain dasabuvir 200mg, ombitasvir 8.33mg, paritaprevir 50mg, and ritonavir 33.33mg and are taken at the same time daily with a meal. Taking the tablets in a fasting state could result in reduced cure and the development of resistance. Swallow the tablets whole and do not consume alcohol within 4 hours of taking Viekira XR. Do not skip or miss doses. Store the medication at or below 86°F.

Do not take more than the prescribed dose of Viekira Pak™ or Viekira XR™ (no doubling up). Do not take Viekira Pak™ or Viekira XR™ if you have had a severe skin rash after taking ritonavir (Norvir®).

The most common side effects of Viekira Pak™ or Viekira XR™ are nausea, itching, and sleep problems.

Tell your healthcare provider if you are taking any of the following medicines; as they are contraindicated with Viekira Pak™ or Viekira XR™:

- Alfuzosin hydrochloride (Uroxatral™)
- Colchicine (COLCRYS, Mitigare®)
- Carbamazepine (Carbatrol™, Eptol™, Equetro™, Tegretol™); Phenytoin (Dilantin™, Phenytek™); Phenobarbital (Luminal™); Primidone (Mysoline®)
- Dronedarone (Multaq®); Lurasidone (Latuda®)
- Efavirenz (Atripla™, Sustiva™)
- Ergot containing medicines including: ergotamine tartrate (Cafergot™, Ergomar™, Ergostat™, Medihaler™); dihydroergotamine mesylate (D.H.E. 45™, Migranal™); methylergonovine (Methergine®); ergonovine (Ergotrate®)
- Ethinyl estradiol-containing medications; combination birth control pills or patches, such as Lo Loestrin™ FE, Norinyl™, Ortho Tri-Cyclen Lo™, Ortho Evra™; hormonal vaginal rings such as NuvaRing®; hormonal replacement therapy medicine Fem HRT™.
- Gemfibrozil (Lopid™); Lovastatin (Advicor™, Altoprev™, Mevacor™); Simvastatin (Simcor™, Vytorin™, Zocor™)

- Midazolam, when taken by mouth; Triazolam (Halcion™)
- Pimozide (Orap™)
- Rifampin (Rifadin™, Rifamate™, Rifater™, Rimactane™)
- Ranolazine (Ranexa®)
- Sildenafil citrate (Revatio™) when taken for pulmonary artery hypertension
- St. John's wort (Hypericum perforatum) or a product that contains St. John's wort

Tell your healthcare provider if you are taking any of the following medicines; as they are not recommended with Viekira Pak™ or Viekira XR™:

- Darunavir (Prezista®) / ritonavir; lopinavir/ ritonavir (Kaletra®); rilpivirine (Edurant®, Complera®, Odefsey®)
- Salmeterol (Serevent, Advair®)

Tell your healthcare provider if you are taking any of the following medicines; as dosage adjustments or monitoring may be recommended:

- Amiodarone (Cordarone®, Nexterone®, Pacerone®); bepridil; disopyramide (Norpace®, Norpace CR®); flecainide (Tambocor™); systemic lidocaine (Xylocaine®); mexiletine; propafenone (Rythmol, Rythmol SR); quinidine (Nuedexta®)
- Valsartan (Diovan®, Exforge®, Entresto™); losartan (Cozaar®, Hyzaar®); candesartan (Atacand®)
- Amlodipine (Norvasc®); Nifedipine (Procardia®, Adalat®); Diltiazem (Cardizem®, Tiazac®); verapamil (Covera-HS®, Calan®, Verelan®)
- Furosemide (Lasix®)
- Ketoconazole; Voriconazole (Vfend®)
- Fluticasone (Inhaled- Arnuity™ Ellipta®, Breo Ellipta®, Flovent®, Advair®; Nasal – Flonase®, Veramyst, Dymista®)
- Atazanavir (Reyataz®)/ ritonavir
- Cyclosporine (Gengraf®, Neoral®, Sandimmune®); tacrolimus (Astagraf XL®, Envarsus XR™, FK506 (common name), Hecoria™, Prograf®)
- Buprenorphine/naloxone (Suboxone®)
- Omeprazole (Prilosec®, Prilosec OTC®)

- Alprazolam (Xanax®)
- Quetiapine (Seroquel®)
- Rosuvastatin (Crestor®); pravastatin (Pravachol®)

PLEASE NOTE:

You must let your medical, mental health, dental providers, and pharmacist(s) know that you are taking Viekira Pak™ or Viekira XR™ prior to starting any new medications. You must let your providers know about any new medications you are prescribed before starting them. This includes vitamins and other supplements.

***Hepatitis C treatment should not cause pain that requires narcotic pain medication.

BENEFITS OF TREATMENT

In most cases, hepatitis C will respond to treatment as determined by a blood test that measures the presence and amount of hepatitis C in the blood. If you have no hepatitis C in your blood 12 weeks **after** the end of treatment, this is called a “sustained virologic response” and means you no longer have hepatitis C. Your chance of achieving a sustained virologic response depends on the hepatitis C genotype, how much hepatitis C virus you have in your blood at the beginning of treatment, any past treatment response, and how much liver damage you have had prior to treatment.

It is possible that you may develop some serious side effects, which will require you to stop the treatment. You may still benefit from treatment even if it does not get rid of your hepatitis C, as it may slow down the disease. You may choose to stop treatment at any time.

In Clinical Trials:

Persons with genotype 1b, without cirrhosis who were treatment-naïve (never treated before) or treatment-experienced (prior treatment failed), given Viekira Pak™ (the components of Viekira XR™) for 12 weeks had a 100% response (cure) rate. Those with compensated (Child Pugh A) cirrhosis who were either treatment naïve or had prior treatment experience had a 100% response rate.

WHOM TO CALL

If you have any questions about your treatment, contact your primary care provider.

TREATMENT AGREEMENT

To receive treatment, please review the following statements and initial beside the responses:

_____ I agree not to drink alcohol or use recreational drugs during the treatment.

_____ I will tell my provider if I have any serious medical conditions (such as heart disease, high blood pressure, diabetes, high cholesterol, rheumatoid arthritis, or drug addiction), or psychiatric conditions (depression, history of suicide attempts, bipolar disorder, or psychosis).

_____ I am willing to visit the clinic and see a provider on a regular schedule for the entire length of the treatment. If I am unable to attend an appointment, I will let my provider know this ahead of time and I will reschedule my appointment.

_____ I understand that my treatment will be stopped if I cannot attend appointments as required to evaluate my health and well-being during treatment and the effectiveness of treatment.

_____ As a female taking Viekira Pak™ or Viekira XR™, I will not get pregnant or breastfeed while on treatment. I understand that my treatment will be stopped if I become pregnant.

_____ Not applicable, I am surgically sterile or post-menopausal.

_____ If I have any problems with the medications or side effects that bother me, I will let my provider or nurse know right away.

_____ I understand that my hepatitis C may not respond to treatment.

_____ I understand that my provider can stop my treatment if the provider feels that stopping it is in the best interest of my health and welfare.

_____ I will do my best to take my medications as prescribed by my provider. If I am unable to do so, I will contact my provider.

_____ I will protect myself and others from hepatitis C by not sharing needles, toothbrushes, razors or nail clippers and covering cuts to prevent blood exposure.

My signature below means that I have read this treatment agreement and/or the meaning of the information has been explained to me. I agree to treatment.

Patient's Name (PLEASE PRINT)	Patient's Signature	Date
--------------------------------------	----------------------------	-------------

Provider's Name (PLEASE PRINT)	Provider's Signature	Date
---------------------------------------	-----------------------------	-------------

**Viekira Pak™ (Ombitasvir/Paritaprevir/Ritonavir & Dasabuvir) or
Viekira XR™ (Ombitasvir/Paritaprevir/Ritonavir/Dasabuvir) Treatment Medications**

_____ You will take **Viekira Pak™**.

Take 2 pink tablets and 1 beige tablet in the morning with a meal. Take 1 beige tablet in the evening with a meal.

The generic name for Viekira Pak™ is ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets, co-packaged for oral use.

Two pink tablets contain ombitasvir 12.5 mg, paritaprevir 75 mg, and ritonavir 50 mg. The two pink tablets are taken at the same time daily (in the morning) with a meal.

The beige tablets contain dasabuvir 250 mg. Take one beige tablet twice daily (in the morning and evening) with a meal.

_____ You will take **Viekira XR™**.

Take 3 pale yellow-colored tablets each morning with a meal.

The generic name for Viekira XR™ is dasabuvir, ombitasvir, paritaprevir, and ritonavir extended release tablets for oral use.

These tablets contain dasabuvir 200mg, ombitasvir 8.33mg, paritaprevir 50mg, and ritonavir 33.33mg.

You get _____ from _____.

Pick up refills on: _____

Call _____ to schedule your family medicine treatment appointments, or if you have any other health concerns.

*****For any emergencies after normal business hours, please go to the Emergency Room. Make sure any healthcare provider you see knows you are on treatment. Carry a list of your medicines with you.**

For more information on managing side effects

visit: <http://www.anthctoday.org/community/hep/patients/index.html>

Click on "Patient Guide- Managing HepC Treatment"

Viekira Pak™ or Viekira XR™ 12 week Lab Tracking Form

General Patient Information

Name: _____
 DOB: ____/____/____
 MRN: _____
 Phone #: _____
 Treatment Start Date: _____

Pre-Treatment Lab Results

HCV RNA: _____
 Genotype: _____ HIV: ____ TSH: ____
 Vit D 25OH: _____ AFP: _____ GFR: _____
 PT/INR: _____ A1C/Glucose: _____

Medication Regimen

Select which formulation is being taken:

-Viekira Pak™ Do not change dose.
 2 pink tablets of ombitasvir, paritaprevir, ritonavir with breakfast.
 1 beige tablet of dasabuvir with breakfast and 1 with dinner.

OR

-Viekira XR™ Do not change dose.
 3 yellow tablets of dasabuvir, ombitasvir, paritaprevir, ritonavir with a meal.

Completed Treatment Week	Lab Date	Hgb	Hct	WBC	PLT	ALT	AST	Alk Phos	Total Bili	Creat/GFR	PHQ-9 (Specified weeks)	HCV RNA (Specified weeks)	Weight (kg)	Pregnancy Test
Pre-Treatment														
Treatment Start Week 0											PHQ-9	HCV RNA		
<i>optional</i>														
<i>optional</i>														
<i>optional</i>														
Week 4												HCV RNA		
<i>optional</i>														
<i>optional</i>														
Week 8														
<i>optional</i>														
<i>optional</i>														
Week 12												HCV RNA		
<i>optional</i>														
3 months post treatment												HCV RNA		

Labs recommended for each follow up visit: CBC, CMP, pregnancy test (females of childbearing age), and HCV RNA as specified.

Please Remember

Give the End of Treatment Letter to the patient at the completion of treatment.

End of Treatment Letter is found in Treatment Monitoring section on webpage.

12 weeks after treatment completion obtain an HCV RNA to check for a sustained virologic response (SVR). SVR is considered a virologic cure of hepatitis C.

SVR12 Cure Letter is found in Treatment Monitoring section on webpage.

<http://anthctoday.org/community/hep/providers/treatment/index.html>