

Commonly Prescribed Hepatitis C Treatment Options and Considerations

(listed in alphabetical order)

Treatment Option Click on medication below for prescribing information	# of pills/day	Treatment Length	What Genotypes ?	Special Considerations Note: Check drug interactions with all Hep-druginteractions.org	Use with Decompensated Cirrhosis?	Use in Severe Renal Disease (GFR < 30)?*
Elbasvir/Grazoprevir (Zepatier™)	1	12-16 weeks	1, 4	If genotype 1a, test for NS5A RAS prior to treatment	No	Yes
Glecaprevir/Pibrentasvir (Mavyret™)	3 at same time with food	8-12 weeks	All	Not to use with ethinyl estradiol containing meds such as combined oral contraceptives	No	Yes
Ledipasvir/Sofosbuvir (Harvoni®)	1	8-12 weeks	1, 4, 5, 6	Acid sensitive. See PI for specific info re: dosing PPIs/acid reducing agents	Yes**	No
Sofosbuvir/Velpatasvir (Epclusa®)	1	12 weeks	All	Acid loving. PPI use not recommended. See PI for specific info re: PPIs/acid reducing agents RAS testing G3 cirrhosis	Yes**	No
Sofosbuvir/Velpatasvir/Voxelaprevir (Vosevi®)	1	12 weeks	All	For use in re-treatment of HCV	No	No

Note: All treatment options had sustained virologic response rates of 94% or better in clinical trials. Side effects commonly seen were headache and fatigue (>10% in clinical trials).

*Consult Liver Disease Specialist before prescribing in severe renal disease.

**Taken with ribavirin. Consult Liver Disease Specialist before prescribing in decompensated cirrhosis.