

Hepatitis C Treatment Medications & Dosing

Please refer to each drug's FULL prescribing information (PI) and see Decision Trees for choosing specific treatment regimen. These can be found on our treatment page: <http://anthc.org/hep>

Daklinza™ (Daclatasvir) with Sofosbuvir with or without Ribavirin

Daklinza™ is a 60mg green pentagonal-shaped tablet, taken once daily with or without food. Dosage change should be discussed with Liver Disease Specialist.

Epclusa® (Sofosbuvir/Velpatasvir) with or without Ribavirin

Epclusa® is a fixed-dose combination, pink diamond-shaped tablet of sofosbuvir 400mg and velpatasvir 100mg, taken once daily with or without food. The dose should not be changed. Velpatasvir requires an acidic environment; refer to PI for recommendations when using in conjunction with H2 antagonists or antacids. Use with PPIs is not recommended.

Harvoni® (Ledipasvir/Sofosbuvir) with or without Ribavirin

Harvoni® is a fixed-dose combination, orange diamond-shaped tablet of ledipasvir 90mg and sofosbuvir 400mg, taken once daily with or without food. The dose should not be changed. Ledipasvir absorption requires an acidic gastric environment; refer to PI for recommendations when using in conjunction with PPI, H2 antagonists, and antacids.

Mavyret™ (Glecaprevir/Pibrentasvir)

Mavyret™ is three pink colored oblong tablets, of fixed-dose combination glecaprevir 100mg and pibrentasvir 40mg (total daily dose 300mg and 120 mg). All three tablets are taken once daily with food.

Olysio® (Simeprevir) with Sofosbuvir

Olysio® is a white 150mg capsule, taken once daily with food. The dose should not be changed.

Sovaldi® (Sofosbuvir) with Ribavirin and/or Daclatasvir, or Simeprevir

Sovaldi® is a yellow capsule-shaped 400mg tablet, taken once daily with or without food. The dose should not be changed. It is prescribed with Daclatasvir with or without ribavirin or Simeprevir.

Technivie™ (Ombitasvir/Paritaprevir/Ritonavir) with Ribavirin

Technivie™ is two pink oblong tablets containing ombitasvir 12.5mg, paritaprevir 75mg, and ritonavir 50mg, taken once daily with a meal. The dose should not be changed.

Viekira Pak™ (Ombitasvir/Paritaprevir/Ritonavir & Dasabuvir) or

Viekira XR™ (Dasabuvir/Ombitasvir/Paritaprevir/Ritonavir) with or without Ribavirin

Viekira Pak® is two pink oblong tablets containing ombitasvir 12.5mg, paritaprevir 75mg, ritonavir 50mg and two beige tablets contain dasabuvir 250mg. Take both pink tablets and one beige tablet in the morning with a meal and one beige tablet in the evening with a meal. The dose should not be changed.

Viekira XR™ is three pale yellow-colored tablets containing dasabuvir 200mg, ombitasvir 8.33mg, paritaprevir 50mg, and ritonavir 33.33mg, taken once daily with a meal.

Vosevi® (Sofosbuvir/Velpatasvir/Voxilaprevir)

Vosevi® is a beige, capsule-shaped fixed dose combination tablet of sofosbuvir 400mg, velpatasvir 100mg and voxilaprevir 100mg, taken once daily with food. The dose should not be

Hepatitis C Treatment Medications & Dosing

changed. Velpatasvir requires an acidic environment; refer to PI for recommendations when using in conjunction with PPIs, H2 antagonists, or antacids.

Zepatier™ (Elbasvir/Grazoprevir) with or without Ribavirin

Zepatier™ is a fixed-dose combination, beige oval-shaped tablet of elbasvir 50mg and grazoprevir 100mg taken once daily with or without food. The dose should not be changed.

Ribavirin

Ribavirin comes in a 200 mg capsule or tablet and should be taken with food.

IMPORTANT: Pregnancy Category X.

Please note: If creatinine clearance is below 50mL/min, ribavirin dose reduction is required. Refer to Copegus® prescribing information for full information about ribavirin dose reduction for renal impairment (Section 2.4). Consult Liver Disease Specialist if any additional concerns. Ribavirin dose is weight based (see tables below) and divided between two doses (morning and evening) with food.

Some patients with cirrhosis and baseline anemia, may need to start low dose ribavirin (600mg/day divided into 2 doses 400 mg in the am and 200 mg in the pm) and increase dose based on tolerance. Please consult Liver Disease Specialist when treating a patient with cirrhosis or if baseline Hgb is less than 12.

Ribavirin weight-based dosing for **Daklinza, Epclusa, Harvoni, Sovaldi, Technivie, or Viekira Pak/XR:**

Weight	Ribavirin Dose
< 75 kg	600 mg qAM, 400 mg qPM (1000 mg daily)
≥ 75 kg	600 mg qAM, 600 mg qPM (1200 mg daily)

Ribavirin weight-based dosing for **Zepatier:**

Weight	Ribavirin Dose
< 66 kg	400 mg qAM, 400 mg qPM (800 mg daily)
66–80 kg	600mg qAM, 400 mg qPM (1000 mg daily)
81-105kg	600 mg qAM, 600 mg qPM (1200 mg daily)
>105 kg	800 mg qAM, 600 mg qPM (1400 mg daily)

Ribavirin Dose Reduction for Anemia

When hemoglobin drops below 10g/dl on treatment, ribavirin dose reduction is recommended. See guidelines:

Hgb (g/dl)	Action
≥ 10	Continue at weight-based dose
< 10	Reduce ribavirin to 600 mg/day divided into 2 doses (400mg am & 200mg pm)
< 8.5	Discontinue ribavirin temporarily and consult liver disease specialist. Once ribavirin has been held, an attempt may be made to restart at 600 mg/day and further increase to 800 mg/day. However, it is not recommended that ribavirin be increased to the original assigned dose (1000 or 1200mg/day).
Patients with history of stable cardiac disease and a decrease in Hgb of ≥ 2 g/dL during any 4-week treatment period, reduce ribavirin to 400 mg every morning and 200 mg every evening; discontinue if Hgb less than 12 g/dL after 4 weeks of reduced dose; if Hgb values improve, may restart at 600 mg daily with further increase to 800 mg daily; do not resume original assigned dose.	