



## ALASKA NATIVE TRIBAL HEALTH CONSORTIUM

Liver Disease & Hepatitis Program  
4315 Diplomacy Drive, Anchorage, AK 99508  
Phone: 907-729-1560 Fax: 907-729-1570  
<http://www.anthc.org/hep>

We are glad to hear you are interested in treatment for HCV! Here are some things to think about (and do) before you make your decision about treatment:

**Why be treated?** Current medications have high cure rates and less side effects. Getting rid of HCV reduces your risk of developing complications and improves your quality of life.

### **What FDA-approved treatments are available?**

These are commonly used treatments for **Genotype 1**:

- Option 1 is Mavyret™ (glecaprevir/pibrentasvir), 3 tablets taken once daily with food for 8-12 weeks. The most common side effects are headache (18%) and fatigue (15%). In clinical studies, the treatment response rate to Mavyret™ was 99% for genotype 1.
- Option 2 is Harvoni® (ledipasvir/sofosbuvir), 1 tablet taken once a day for 8-12 weeks. The most common side effects are feeling tired (16%) and headache (14%). In clinical studies, treatment response rates to Harvoni® were 94-100%.
- Option 3 is Epclusa® (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The most common side effects are headache (22%) and feeling tired (15%). In clinical studies, treatment response rates to Epclusa® were 94-98% for genotype 1.

These are commonly used treatment options for **Genotype 2**:

- Option 1 is Mavyret™ (glecaprevir/pibrentasvir), 3 tablets taken once daily with food for 8-12 weeks. The most common side effects are headache (18%) and fatigue (15%). In clinical studies, treatment response rates to Mavyret™ were 98-100% for genotype 2.
- Option 2 is Epclusa® (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The most common side effects are headache (22%) and feeling tired (15%). In clinical studies, the treatment response rate to Epclusa® was 99% for genotype 2.

These are commonly used treatment options for **Genotype 3**:

- Option 1 is Mavyret™ (glecaprevir/pibrentasvir), 3 tablets taken once daily with food for 8-12 weeks. The most common side effects are headache (18%) and fatigue (15%). In clinical studies, treatment response rates to Mavyret™ were 95-98% for genotype 3.
- Option 2 is Epclusa® (sofosbuvir/velpatasvir), 1 tablet taken once a day for 12 weeks. The most common side effects are headache (22%) and feeling tired (15%). In clinical studies, treatment response rates to Epclusa® were 85-98% for genotype 3.

There are no data on the new HCV drugs in pregnant women or nursing mothers. Safety/risk during pregnancy or breastfeeding has not been established.

Some treatments will require ribavirin which is 5-6 more tablets per day divided between morning and evening with food. The major side effects are feeling tired, nausea, itching and skin rash, trouble sleeping, irritability and weakness. A common side effect of ribavirin is anemia. **PLEASE NOTE: Ribavirin cannot be given to a pregnant or breastfeeding female or to a female who plans to become pregnant or a male who plans to father a child during or for 6 months after treatment because it can cause birth defects.**

### **Are you ready for treatment?**

To ensure your success in completing HCV treatment, please consider the following:

- Drinking alcohol or misusing opioids or other drugs can hurt the liver. If you have recent drug/alcohol abuse, it is recommended that you attend an approved drug treatment program for support.
- Have you discussed HCV treatment with your primary care provider?
- It's helpful to have a relative/close friend to support you during treatment.
- Are you committed to making every treatment appointment and getting **monthly** blood draws? We will want to monitor your progress during treatment.
- Can you return for an appointment 12 weeks after completing medication to be tested for cure?

### **Additional Requirements If Checked:**

If you have cirrhosis, you may need an:

\_\_\_\_ EGD (A tube put into the esophagus and stomach to look for swollen veins that can bleed).

\_\_\_\_ Ultrasound of the liver (done in the past 6 months). This non-invasive test checks your liver for cancer.

Call your primary care provider to make an appointment to discuss HCV treatment once you are ready.

If you are coming to Anchorage and want a FibroScan, call the Liver Clinic ahead of your visit to schedule an appointment. FibroScan is a test using ultrasound waves to check liver stiffness or scarring/fibrosis in your liver. FibroScan testing is done in the Internal Medicine Clinic. Do not eat or drink for 3 hours before the test. Call us at 907-729-1560 or 1-800-655-4837.