Bisacodyl/Magnesium Citrate Split Prep

How to prepare for your colonoscopy

QUESTIONS? Call 907-729-4444 to speak with a Colorectal Screening Clinic staff member. We are here to help you!

Variation and the state of the distant	_		
Your procedure is scheduled for:	Date	Time	PROVIDER NOTES:
	Mon Tue	Wed Thu Fri	
Check-in at the Ambulatory SurgerDrive, first floor and expect to stay OR	•		sity Lake
Check-in to ANMC Day Surgery, lo floor of the main hospital for your		•	
 AT LEAST FIVE DAYS BEFORE YOUR PR Stop taking any iron supplements a any blood thinners should be stopp Please call Central Registration at 90 procedure to pre-register. If you live in-person to register if you are able 	nd discuss woed until afte 07-729-1395 ue in Anchora	er the procedure. up to one week befo	ore your
 AT LEAST ONE DAY BEFORE YOUR PRO Go to Pharmacy to pick up your bowhich will be two – 10 oz. (296ml) be Refrigerate bottles ahead of time to 	wel preparat ottles of ma	gnesium citrate solu	ition.
THE DAY BEFORE YOUR PROCEDURE (or the Friday	, before a Monday p	orocedure):

One day before your procedure start a clear liquid diet and no solid foods.

A Day Surgery nurse will be calling you to give you the check-in time for the day of surgery. Please make sure your phone is turned on and able to receive phone calls.

If you do not get a call by 6 p.m., please call 907-729-1156 for assistance.

Date: Time:



Food and drink you cannot have:

- x Nuts or seeds
- x Alcoholic beverages
- X Ice cream or smoothies
- x Foods like milk or cream
- X Breads, meats, eggs, or dairy
- x Soy milk, almond milk or rice milk
- x Juices that are red/orange or have pulp like orange, grapefruit, or tomato juice









Clear liquids you can eat or drink

- Water, sparkling water, and/or clear flavored water
- Black coffee and/or tea with sugar and/or honey (No milk, no creamer)
- Clear chicken, beef, or vegetable STOCK OR BROTH ONLY! (No pieces of meat, eggs or noodles in broth)
- Jell-O or popsicles (No red color)
- Clear fruit juices like apple or white grape (No pulp and no red color)
- 7-Up, Sprite, Gatorade, Crystal Light, ginger-ale, or lemonade (No red color)

Bisacodyl/Magnesium Citrate Split Prep

How to take the medication



You must complete the entire prep to ensure the most effective colonoscopy.

One day before the procedure take Dose 1

At 4 p.m.

Take the 4 bisacodyl (Dulcolax) tablets.

At 5 p.m.

Drink Magnesium Citrate according to the following instructions.

- Drink 1 container of Magnesium Citrate (10oz or 296ml). Refrigerate the bottle ahead of time to improve the taste.
- Immediately after drinking Magnesium Citrate, drink at least 3 cups of clear liquids.
- Continue to drink a lot of CLEAR liquids up to 5:30 a.m. the day of your procedure. The more clear liquids you drink, the better the prep will be.

After Midnight

- Nothing to smoke
- No chewing tobacco

The day of your procedure take Dose 2

At 4:30 a.m.

Drink 1 container of Magnesium Citrate (10oz or 296ml). Refrigerate the bottle ahead of time to improve the taste.

- Immediately after drinking Magnesium Citrate, drink at least 3 cups of clear liquids.
- You may drink clear liquids, including black coffee without cream, up to 5:30 a.m. the day of your procedure.

You are done with the prep.Good job, the hardest part is over!

TIPS FOR PROPER HYDRATION

Prevent fluid loss and dehydration by drinking clear liquids before, during, and after taking the Magnesium Citrate solution. It is important for you to drink an additional 3-4 cups of clear liquids after Dose 1 and Dose 2 to ensure the most effective cleansing without becoming dehydrated.

Remember to stop drinking clear liquids at 5:30 a.m. the morning of your procedure.

TIP: Using baby wipes and/or Tucks® medicated pads may help with anal soreness during bowel preparation.

Take all medications as normal unless otherwise instructed. Remember to stop taking any blood thinners as instructed.



REMINDER!

No Driving Permitted after your procedure.

You must have a responsible escort who is 18 or older who can take you home and stay with you for 24 hours after your procedure.

After your procedure, it is normal to feel bloating or mild abdominal cramping. Passing gas will help you feel better.

If you have a fever or chills, significant bleeding or severe abdominal pain, contact us at 907-729-4444.

If a biopsy was done or polyps were removed, results usually take at least 2 weeks to obtain but sometimes can take longer.