HCV Simplified Treatment
For Alaska Tribal Health System

Updated 8/2024
For more information, visit:
www.anthc.org/hep

Start here:

Step 1

Calculate FIB-4
https://www.hepatitisc.uw.edu/page/clinical-calculators/fib-4

and

Assess for Cirrhosis:
- FIB-4 > 3.25 or any of the following:
- Transient elastography (e.g. FibroScan stiffness > 12.5 kPa
- Noninvasive serologic tests above proprietary cutoffs for cirrhosis (e.g. FibroSure, FibroTest)
- Clinical evidence of cirrhosis (liver nodularity and/or splenomegaly on imaging, platelet count < 150,000
- Prior liver biopsy showing cirrhosis

FIB-4 < 3.25 and no clinical signs of cirrhosis

FIB-4 > 3.25 or clinical signs of cirrhosis

Calculate CTP score
https://www.hepatitisc.uw.edu/page/clinical-calculators/ctp

CTP ≤ 6

CTP > 6

Proceed with Simplified Treatment Step 2

Note: Patients with compensated cirrhosis can be treated in Primary Care setting. For questions about cirrhosis and treatment, please consult Liver Clinic or refer for evaluation. Persons with cirrhosis should have annual follow up with Liver Disease specialist.

Refer to Liver Clinic or a medical practitioner with expertise in liver disease